

## LETTERS

Author affiliations: Etablissement Français du Sang Alpes-Méditerranée, Marseille, France (P. Biagini, M. Touinssi, C. Picard, P. Gallian, J.-F. Cantaloube, P. de Micco); Université de la Méditerranée, Marseille (P. Biagini, M. Touinssi, P. Gallian, J.-F. Cantaloube, P. de Micco); Centre Hospitalier Universitaire, Marseille (B. Dussol); Hôpital de la Conception, Assistance Publique de Marseille, Marseille (P. Brunet; V. Moal, Y. Berland); and Institute for Animal Health, Pirbright, UK (M. Belhouchet, H. Attoui)

DOI: 10.3201/eid1411.080862

## References

1. Jones MS, Kapoor A, Lukashov VV, Simmonds P, Hecht F, Delwart E. New DNA viruses identified in patients with acute viral infection syndrome. *J Virol*. 2005;79:8230–6. DOI: 10.1128/JVI.79.13.8230-8236.2005
2. Fryer JF, Delwart E, Bernardin F, Tuke PW, Lukashov VV, Baylis SA. Analysis of two human parvovirus PARV4 genotypes identified in human plasma for fractionation. *J Gen Virol*. 2007;88:2162–7. DOI: 10.1099/vir.0.82620-0
3. Fryer JF, Kapoor A, Minor PD, Delwart E, Baylis SA. Novel parvovirus and related variant in human plasma. *Emerg Infect Dis*. 2006;12:151–4.
4. Fryer JF, Lucas SB, Padley D, Baylis SA. Parvoviruses PARV4/5 in hepatitis C virus-infected patient. *Emerg Infect Dis*. 2007;13:175–6.
5. Fryer JF, Delwart E, Hecht FM, Bernardin F, Jones MS, Shah N, et al. Frequent detection of the parvoviruses, PARV4 and PARV5, in plasma from blood donors and symptomatic individuals. *Transfusion*. 2007;47:1054–61. DOI: 10.1111/j.1537-2995.2007.01235.x
6. Fryer JF, Hubbard AR, Baylis SA. Human parvovirus PARV4 in clotting factor VIII concentrates. *Vox Sang*. 2007;93:341–7.
7. Longhi E, Bestetti G, Acquaviva V, Foschi A, Piolini R, Meroni L, et al. Human parvovirus in the bone marrow of Italian patients with AIDS. *AIDS*. 2007;21:1481–3. DOI: 10.1097/QAD.0b013e3281e38558
8. Manning A, Willey SJ, Bell JE, Simmonds P. Comparison of tissue distribution, persistence, and molecular epidemiology of parvovirus B19 and novel human parvoviruses PARV4 and human bocavirus. *J Infect Dis*. 2007;195:1345–52. DOI: 10.1086/513280
9. Schneider B, Fryer JF, Reber U, Fischer HP, Tolba RH, Baylis SA, et al. Persistence of novel human parvovirus PARV4 in liver tissue of adults. *J Med Virol*. 2008;80:345–51. DOI: 10.1002/jmv.21069

Address for correspondence: Philippe Biagini, UMR CNRS 6578 Equipe “Emergence et co-évolution virale,” Etablissement Français du Sang Alpes-Méditerranée et Université de la Méditerranée, 27 Blvd Jean Moulin, 13005 Marseille, France; email: [philippe.biagini@univmed.fr](mailto:philippe.biagini@univmed.fr)

## Establishment of *Biomphalaria tenagophila* Snails in Europe

**To the Editor:** Schistosomiasis, known since ancient times, is caused by blood flukes (Trematoda: Schistosomidae). It is a major communicable disease with public health and socioeconomic effects in the developing world (1). Among parasitic diseases, schistosomiasis ranks second only to malaria with regard to the number of persons infected and at risk. The life cycle of schistosomes is complex, requiring specific freshwater snails as intermediate hosts for larvae development and multiplication. Among *Schistosoma* species that affect humans, *Schistosoma mansoni* is the most likely to invade new areas mainly because of the adaptability and invasiveness of its intermediate host, *Biomphalaria* snails. Natural populations of these snails are usually found in tropical standing water or freshwater in South America and Africa, but they also reach 30° latitude in subtropical areas (1,2). Many species of these red-blooded planorbid snails (Gastropoda: Basommatophora) are able to survive a long time when removed from their freshwater habitat (1). Of the 34 *Biomphalaria* species, 4 (*B. glabrata*, *B. pfeifferi*, *B.*

*straminea*, and *B. tenagophila*) have recently expanded their native ranges (3). They have been introduced to areas where other *Biomphalaria* species are endemic (e.g., Congo and Egypt) or to subtropical zones that have no frost period (Texas, Louisiana, Florida, Hong Kong) (3,4). None of the known invasions, whether peripheral range expansion or long distance dispersal, reached the temperate zone. Spreading of the blood-fluke snails to schistosome-free areas may enable the parasite to colonize new habitats concurrently, expanding the potential area of clinical schistosomiasis.

We collected these snails in spring 2005, autumn 2006, and autumn 2007, near Răbăgani, Romania, Eastern Europe (46°45'1.3"N, 22°12'44.8"E) in a hypothermal spring. Water temperature was 25°C in the spring and 16°C–25°C, gradually decreasing, along the brook course. In and beside an abandoned concrete pool next to the spring, we collected 100 shells and 34 living specimens that macroscopically resembled *Biomphalaria* spp. snails. All 16 dissected animals proved to be fully developed adults, according to the maturity of their genital organs (Figure). Using available identification keys (5), we tentatively identified these snails as *B. tenagophila*. Voucher specimens have been deposited in the Hungarian Natural History Museum (accession nos. HNHN96857 and HNHN95433).

DNA was extracted from the foot muscles of 3 specimens by using QIAamp DNA Mini Kit (QIAGEN, Hilden, Germany). For amplification of the partial mitochondrial 16S ribosomal RNA gene, we used a PCR with primers 16Sar and 16Sbr (6). Nucleotide sequences were determined in both directions. PCR products of ≈430 bp were detected from all 3 samples. Automatic cycle sequencing of the randomly selected amplicon (GenBank accession no. EU069412) showed 99.74% similarity to *B. tenagophila* (AF449615, Brazil).

Our morphologic, anatomic, and molecular data unambiguously prove the occurrence of *B. tenagophila* snails in Romania. *B. tenagophila* snails had been found earlier (in 2004) at this location but had presumably been misidentified as dwarf specimens of a common European species, *Planorbis corneus* (7). Consequently, *B. tenagophila* snails have been not only introduced, but also established in Răbăgani, representing the furthest self-sustaining population of this species from the equator.

*B. tenagophila* is a new species for the European fauna. It could represent a founder population of unknown origin for further spread into Europe, which might easily be accomplished by migrating birds or more likely by plants used in aquariums (3). Although no trematode larvae were detected in the observed specimens, clinical schistosomiasis can be imported by immigrants or tourists into Europe, as has been reported in Romania and neighboring Hungary (8,9). If eggs were released in feces of humans infected with the blood flukes, they could hatch in the environment and the larvae

could develop to an infective stage in these snails. The observed local cultural and social factors involving natural water (washing clothes, bathing) in Răbăgani where *B. tenagophila* have been found may also increase the chance of human infection.

We believe that *B. tenagophila* in Europe, together with the global climate change and a possible encounter of these snails with schistosomes, could pose a public health risk. Measures must be taken to prevent the spread of this species into European freshwater. Chemical control is not possible in Răbăgani because it is an area where other rare and endangered snail species are protected (7). Therefore, the manual collection and removal of all the *B. tenagophila* specimens in the area seems to be the only possibility for eradication, which might remain in effect for years. To avoid similar establishments, we suggest regular malacologic and parasitologic surveillance of at least the thermal and hypothermal water bodies for these tropical invaders around European settlements.

## Acknowledgments

We thank W.M. Lotfy, V. Estrada, R.L. Caldeira, and M.J. Kenny for their comments.

This work was partly supported by the Hungarian National R&D Program, "The origin and genesis of the fauna of the Carpathian Basin: diversity, biogeographical hotspots and nature conservation significance; contract no. 3B023-04."

**Gábor Majoros,<sup>1</sup> Zoltán Fehér, Tamás Deli, and Gábor Földvári<sup>1</sup>**

Author affiliations: Szent István University Faculty of Veterinary Science, Budapest, Hungary (G. Majoros, G. Földvári); Hungarian Natural History Museum, Budapest (Z. Fehér); and Munkácsy Mihály Museum, Békéscsaba, Hungary (T. Deli)

DOI: 10.3201/eid1411.080479

## References

1. Malek EA. Snail-transmitted parasitic diseases, vol. I. Boca Raton (FL): CRC Press, Inc.; 1980.

<sup>1</sup>These authors contributed equally to this article.




Figure. A) Shell morphology of *Biomphalaria tenagophila* snail from Romania. Diameter of the shell was 10–14 mm. The sinistrally coiled, flat shells are yellow-brown, discoidal, deeply and symmetrically biconcave, and consist of 5 or 6 slowly increasing whorls. The last whorl is rounded; the intermediate whorls are slightly angled on the left side. The aperture is circular or slightly ovate and angled toward the left side of the shell (i.e., toward the upper surface on the bottom right shell). Fine, parallel, rib-like transverse lines can be seen on the outer surface of the whorls. A series of photographs were prepared by focusing on different levels of the structure and these were combined by CombineZ5 ([www.hadleyweb.pwp.blueyonder.co.uk](http://www.hadleyweb.pwp.blueyonder.co.uk)), using "do combine" and "do average and filter" commands. Scale bar = 3 mm. B) Reproductive system of *B. tenagophila* snail from Romania; ag, albumin gland; bc, bursa copulatrix; hp, distal part of the hepatopancreas; ng, nidamental gland; od, oviduct; ot, ovotestis; pp, preputium; pr, prostate; ps, penis sheath; sd, spermiduct; sv, seminal vesicles; vd, vas deferens; vp, vaginal pouch. *B. tenagophila*-specific characteristics (5): >200 diverticulae of the ovotestis; 7–11 main lobes of the prostate; and presence of vaginal pouch. Scale bar = 1 mm.

2. Paraense WL. The schistosome vectors in the Americas. Mem Inst Oswaldo Cruz. 2001;96(Suppl):7–16. DOI: 10.1590/S0074-02762001000900002
3. Pointier JP, David P, Jarne P. Biological invasions: the case of planorbid snails. J Helminthol. 2005;79:249–56. DOI: 10.1079/JOH2005292
4. Malek EA. Studies on “tropicorbid” snails (Biomphalaria: Planorbidae) from the Caribbean and Gulf of Mexico areas, including the Southern United States. Malacologia. 1969;7:183–209.
5. Malek EA. Snail hosts of schistosomiasis and other snail-transmitted diseases in tropical America: a manual. Scientific publication 478. Washington: Pan American Health Organization; 1985.
6. DeJong RJ, Morgan JA, Wilson WD, Al-Jaser MH, Appleton CC, Coulbaly G, et al. Phylogeography of *Biomphalaria glabrata* and *B. pfeifferi*, important intermediate hosts of *Schistosoma mansoni* in the New and Old World tropics. Mol Ecol. 2003;12:3041–56. DOI: 10.1046/j.1365-294X.2003.01977.x
7. Gagiú A. On the extinction of the relict snail *Theodoxus prevostianus* Pfeiffer 1828 in Răbăgani, Romania. Nymphaea. 2004;31:75–81.
8. Dancescu P, Colea A, Gafiteanu L, Sima A. Imported tropical parasitosis in Rumania [in French]. Bull Soc Pathol Exot Filiales. 1976;69:501–7.
9. Dobi S, Sántha M, Horváth A, Kassas AL, Várnai F. Introduced schistosomiasis with late complications [in Hungarian]. Orv Hetil. 1995;136:1673–5.

Address for correspondence: Gábor Földvári, Department of Parasitology and Zoology, Faculty of Veterinary Science, Szent István University, 2nd István St, Budapest H-1078, Hungary; email: foldvarigabor@gmx.de

### Letters

Letters commenting on recent articles as well as letters reporting cases, outbreaks, or original research are welcome. Letters commenting on articles should contain no more than 300 words and 5 references; they are more likely to be published if submitted within 4 weeks of the original article's publication. Letters reporting cases, outbreaks, or original research should contain no more than 800 words and 10 references. They may have 1 Figure or Table and should not be divided into sections. All letters should contain material not previously published and include a word count.

## *Rickettsia aeshlimannii* Infection, Algeria

**To the Editor:** Only 2 cases of *Rickettsia aeshlimannii* infection have been reported. We report 2 additional cases documented in Algeria by immunofluorescence (IF) assays and confirmed by Western blot (WB) assays and cross-adsorption studies.

Tick-borne rickettsioses are now recognized as emerging or reemerging human infections worldwide. These zoonoses, caused by intracellular bacteria within spotted fever group (SFG) *Rickettsia* spp., share characteristic clinical features including fever, rash, and sometimes inoculation eschar at the bite site (1). In North Africa, cases of rickettsioses are rarely documented (2). In Algeria, only Mediterranean spotted fever caused by *R. conorii* has been described (3).

From 2000 through 2006 in Algeria, all patients with suspected rickettsioses seen at the infectious diseases units of Constantine and Batna hospitals were included in a prospective study; clinical and epidemiologic data and acute-and convalescent-phase serum samples obtained 2–4 weeks later were collected. Serum samples were sent to Marseille, France, where they were analyzed by an IF assay, using 9 SFG rickettsial antigens (*R. conorii conorii*, *R. conorii israelensis*, *R. africae*, *R. sibirica mongolitimonae*, *R. aeshlimannii*, *R. massiliae*, *R. helvetica*, *R. slovaca*, and *R. felis*) and a typhus group antigen (*R. typhi*) (3). The IF assay result was considered positive 1) if immunoglobulin (Ig) G titers were  $\geq 128$  and/or IgM titers were  $\geq 64$  for *R. conorii* and 2) if IgG titers were  $\geq 64$  and/or IgM titers were  $\geq 32$  for other rickettsial antigens (3). When cross-reactions between several antigens were noted, rickettsial antigen was considered to represent the infectious agent if titers of IgG and/or IgM antibody against this antigen were at least 2-fold

higher than titers of IgG and/or IgM antibody against other rickettsial antigens (3,4). When the difference in titers among several antigens was lower than 2-fold, WB assays and cross-adsorption studies were performed (4,5). A total of 135 patients were included in the study. We describe 2 cases of *R. aeshlimannii* infection. Cases caused by other SFG rickettsiae will be reported elsewhere.

An 80-year-old man who reported contact with dogs parasitized by ticks had a 7-day history of high fever, headache, myalgia, and vomiting. On physical examination, a generalized maculopapular rash, 2 eschars (right shoulder and knee), and bilateral hemorrhagic signs on the retina were noticed. Elevated levels of liver enzymes (aspartate aminotransferase 187 U/L, alanine aminotransferase 108 U/L), hyponatremia (sodium 120 mmol/L), and hypokalemia (potassium 2.9 mmol/L) were found. IF assay showed raised levels of IgG/IgM against *R. aeshlimannii* (512/64) and *R. conorii* (128/0).

The second patient, a 36-year-old man, reported a 15-day history of fever with headache and failure of amoxicillin and cotrimoxazole treatments. Oral aphthous, a maculopapular rash, and purpuric lesions on the arms were noticed. IF assay showed raised levels of IgG/IgM at the same titer (2,048/32) against *R. conorii*, *R. aeshlimannii*, and *R. massiliae*. WB assays and cross-adsorption studies confirmed that antibodies were directed against *R. aeshlimannii* (Figure). Both patients recovered after doxycycline treatment (1).

*R. aeshlimannii* was first characterized as a new SFG rickettsia after its isolation from *Hyalomma marginatum marginatum* ticks in Morocco in 1997 (6). Thereafter, *R. aeshlimannii* has been detected in this tick species in southern Europe and North Africa (7), as well as in *H. m. rufipes* in sub-Saharan Africa (1). Preliminary data have suggested that these *Hyalomma*